

# Dominating Sequences in Graphs

Didem Gözüpek

Department of Computer Engineering, Gebze Technical University, Turkey

CIMPA 2023

# Outline

- 1 Introduction
- 2 Characterization of  $k$ -Uniform Graphs for  $k \leq 3$
- 3 Characterization of  $k$ -Uniform Graphs for  $k \geq 4$




## Preliminaries

- Let  $G = (V(G), E(G))$  be a simple graph with vertex set  $V(G)$  and edge set  $E(G)$
- The *(open) neighborhood* of a vertex  $v \in V(G)$ , denoted by  $N(v)$ , is the set of vertices adjacent to  $v$
- $N[v] = N(v) \cup \{v\}$  = The *closed neighborhood* of a vertex  $v \in V(G)$
- The degree of a vertex  $u$  is  $\deg(u) = |N(u)|$  and minimum degree is denoted by  $\delta(G)$

# Dominating Set

A subset  $S \subseteq V(G)$  is called a *dominating set* of  $G$  if every vertex in  $V(G) \setminus S$  has at least one neighbor in  $S$

$\gamma(G)$  = Domination number = Size of a minimum dominating set



# Dominating Sequences

A sequence  $S = (v_1, \dots, v_k)$  of distinct vertices in a graph  $G$  is called a *legal dominating sequence* if every vertex in the sequence dominates at least one vertex not dominated by those vertices that precede it, and at the end all vertices of  $G$  are dominated; that is, if

$$N[v_i] \setminus \bigcup_{j=1}^{i-1} N[v_j] \neq \emptyset \text{ holds for every } i \in \{2, \dots, k\}.$$

## Dominating Sequences

A sequence  $S = (v_1, \dots, v_k)$  of distinct vertices in a graph  $G$  is called a *legal dominating sequence* if every vertex in the sequence dominates at least one vertex not dominated by those vertices that precede it, and at the end all vertices of  $G$  are dominated; that is, if

$$N[v_i] \setminus \bigcup_{j=1}^{i-1} N[v_j] \neq \emptyset \text{ holds for every } i \in \{2, \dots, k\}.$$

Clearly, the length  $k$  of a legal dominating sequence  $S = (v_1, \dots, v_k)$  is bounded from below by the domination number  $\gamma(G)$ .

## Grundy Domination Number

Bresar et.al. introduced the notion of *Grundy domination number*  $\gamma_{gr}(G)$  as the maximum length of a legal dominating sequence.

The corresponding sequence is called a *Grundy dominating sequence*

## Grundy Domination Number

Bresar et.al. introduced the notion of *Grundy domination number*  $\gamma_{gr}(G)$  as the maximum length of a legal dominating sequence.

The corresponding sequence is called a *Grundy dominating sequence*

The lower bound  $\gamma_{gr}(G) \geq \gamma(G)$  is obvious.



## Grundy Domination Number

Bresar et.al. introduced the notion of *Grundy domination number*  $\gamma_{gr}(G)$  as the maximum length of a legal dominating sequence.

The corresponding sequence is called a *Grundy dominating sequence*

The lower bound  $\gamma_{gr}(G) \geq \gamma(G)$  is obvious.

### Proposition

For an arbitrary graph  $G$ ,  $\gamma_{gr}(G) \leq |V(G)| - \delta(G)$

# Grundy Domination Number

## Proof.

Let  $S = (s_1, \dots, s_k)$  be a Grundy dominating sequence of  $G$  and let  $u$  be a vertex footprinted in the last step. Since  $u$  is not dominated before the last step, we have:

$N[u] \cap \{s_1, \dots, s_{k-1}\} = \emptyset$ , and so

# Grundy Domination Number

## Proof.

Let  $S = (s_1, \dots, s_k)$  be a Grundy dominating sequence of  $G$  and let  $u$  be a vertex footprinted in the last step. Since  $u$  is not dominated before the last step, we have:

$N[u] \cap \{s_1, \dots, s_{k-1}\} = \emptyset$ , and so

$$|\{s_1, \dots, s_{k-1}\}| = k - 1 \leq |V(G)| - (\deg(u) + 1).$$

Thus,  $\gamma_{gr}(G) = k \leq |V(G)| - \delta(G)$



## $k$ -Uniform Graphs

If all legal dominating sequences of a graph  $G$  have the same length  $k$ , that is,  $\gamma(G) = \gamma_{gr}(G) = k$ , then such a graph is called a  *$k$ -uniform graph*.

## $k$ -Uniform Graphs

If all legal dominating sequences of a graph  $G$  have the same length  $k$ , that is,  $\gamma(G) = \gamma_{gr}(G) = k$ , then such a graph is called a  *$k$ -uniform graph*.

Two distinct nodes  $x$  and  $y$  in a graph are called *twins* if  $N[x] = N[y]$ .

## $k$ -Uniform Graphs

If all legal dominating sequences of a graph  $G$  have the same length  $k$ , that is,  $\gamma(G) = \gamma_{gr}(G) = k$ , then such a graph is called a  *$k$ -uniform* graph.

Two distinct nodes  $x$  and  $y$  in a graph are called *twins* if  $N[x] = N[y]$ .

A graph  $G$  is said to be *twin-free* if no two of its distinct nodes are twins.

## $k$ -Uniform Graphs

If all legal dominating sequences of a graph  $G$  have the same length  $k$ , that is,  $\gamma(G) = \gamma_{gr}(G) = k$ , then such a graph is called a  $k$ -uniform graph.

Two distinct nodes  $x$  and  $y$  in a graph are called *twins* if  $N[x] = N[y]$ .

A graph  $G$  is said to be *twin-free* if no two of its distinct nodes are twins.

Creating a twin of a vertex changes neither the domination number nor the Grundy domination number

Therefore, the question of characterizing  $k$ -uniform graphs is interesting only for twin-free graphs

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

We will review the following work:

B.Bresar, T.Gologranc, M.Milanic, D.F.Rall, R.Rizzi, "Dominating sequences in graphs", *Discrete Mathematics*, vol.336, pp.22-36, 2014.



## Characterization of $k$ -Uniform Graphs

### Lemma

*Let  $G$  be a twin-free  $k$ -uniform graph and let  $x, y \in V(G)$ . If  $N[x] \subseteq N[y]$ , then  $x = y$ .*

## Characterization of $k$ -Uniform Graphs

### Lemma

*Let  $G$  be a twin-free  $k$ -uniform graph and let  $x, y \in V(G)$ . If  $N[x] \subseteq N[y]$ , then  $x = y$ .*

### Proof.

Suppose that  $N[x] \subseteq N[y]$ . If  $N[x] = N[y]$ , then we must have  $x = y$  since  $G$  is twin-free.

## Characterization of $k$ -Uniform Graphs

### Lemma

*Let  $G$  be a twin-free  $k$ -uniform graph and let  $x, y \in V(G)$ . If  $N[x] \subseteq N[y]$ , then  $x = y$ .*

### Proof.

Suppose that  $N[x] \subseteq N[y]$ . If  $N[x] = N[y]$ , then we must have  $x = y$  since  $G$  is twin-free. Hence, we may assume that there exists a vertex  $z \in N(y) \setminus N(x)$ .

Since  $G$  is  $k$ -uniform for some  $k$ , the sequence  $S = (x, y)$  can be extended to a legal dominating sequence  $S'$  of  $G$  of length  $k$ , say  $S' = (x, y, v_3, \dots, v_k)$ .

## Characterization of $k$ -Uniform Graphs

### Lemma

*Let  $G$  be a twin-free  $k$ -uniform graph and let  $x, y \in V(G)$ . If  $N[x] \subseteq N[y]$ , then  $x = y$ .*

### Proof.

Suppose that  $N[x] \subseteq N[y]$ . If  $N[x] = N[y]$ , then we must have  $x = y$  since  $G$  is twin-free. Hence, we may assume that there exists a vertex  $z \in N(y) \setminus N(x)$ .

Since  $G$  is  $k$ -uniform for some  $k$ , the sequence  $S = (x, y)$  can be extended to a legal dominating sequence  $S'$  of  $G$  of length  $k$ , say  $S' = (x, y, v_3, \dots, v_k)$ . But then,  $(y, v_3, \dots, v_k)$  is a legal dominating sequence of  $G$  of length  $k - 1$ , contradicting the fact that  $G$  is  $k$ -uniform. □

## Characterization of $k$ -Uniform Graphs

The complement  $\overline{G}$  of a graph  $G$  is a graph on the same vertices such that two distinct vertices of  $\overline{G}$  are adjacent if and only if they are not adjacent in  $G$ .

## Characterization of $k$ -Uniform Graphs

The complement  $\overline{G}$  of a graph  $G$  is a graph on the same vertices such that two distinct vertices of  $\overline{G}$  are adjacent if and only if they are not adjacent in  $G$ .

$G + H$  denotes the disjoint union of  $G$  and  $H$ , and  $nG$  denotes the disjoint union of  $n$  copies of  $G$ .

## Characterization of $k$ -Uniform Graphs

The complement  $\overline{G}$  of a graph  $G$  is a graph on the same vertices such that two distinct vertices of  $\overline{G}$  are adjacent if and only if they are not adjacent in  $G$ .

$G + H$  denotes the disjoint union of  $G$  and  $H$ , and  $nG$  denotes the disjoint union of  $n$  copies of  $G$ .

Let  $G = (V, E)$  be any graph and  $S \subseteq V$ . Then the induced subgraph  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .

## Characterization of $k$ -Uniform Graphs

The complement  $\overline{G}$  of a graph  $G$  is a graph on the same vertices such that two distinct vertices of  $\overline{G}$  are adjacent if and only if they are not adjacent in  $G$ .

$G + H$  denotes the disjoint union of  $G$  and  $H$ , and  $nG$  denotes the disjoint union of  $n$  copies of  $G$ .

Let  $G = (V, E)$  be any graph and  $S \subseteq V$ . Then the induced subgraph  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .

A graph is  $H$ -free if it does not have the graph  $H$  as an induced subgraph.



## Characterization of $k$ -Uniform Graphs

The complement  $\overline{G}$  of a graph  $G$  is a graph on the same vertices such that two distinct vertices of  $\overline{G}$  are adjacent if and only if they are not adjacent in  $G$ .

$G + H$  denotes the disjoint union of  $G$  and  $H$ , and  $nG$  denotes the disjoint union of  $n$  copies of  $G$ .

Let  $G = (V, E)$  be any graph and  $S \subseteq V$ . Then the induced subgraph  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .

A graph is  $H$ -free if it does not have the graph  $H$  as an induced subgraph.

$P_n$  is a path on  $n$  vertices

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- $G$  is 1-uniform if and only if  $G$  is a complete graph*

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- *$G$  is 1-uniform if and only if  $G$  is a complete graph*
- *$G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs*

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- *$G$  is 1-uniform if and only if  $G$  is a complete graph*
- *$G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs*
- *$G$  is 3-uniform if and only if  $G$  is the disjoint union of a 1-uniform and a 2-uniform graph.*

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- *$G$  is 1-uniform if and only if  $G$  is a complete graph*
- *$G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs*
- *$G$  is 3-uniform if and only if  $G$  is the disjoint union of a 1-uniform and a 2-uniform graph.*

An *independent set* is a set of vertices in a graph, no two of which are adjacent.

$\alpha(G)$  denotes the maximum size of an independent set.

### Proof.

$(\Leftarrow)$  Easy.

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

If  $G$  is a graph, then

- $G$  is 1-uniform if and only if  $G$  is a complete graph
- $G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs
- $G$  is 3-uniform if and only if  $G$  is the disjoint union of a 1-uniform and a 2-uniform graph.

An *independent set* is a set of vertices in a graph, no two of which are adjacent.

$\alpha(G)$  denotes the maximum size of an independent set.

### Proof.

( $\Leftarrow$ ) Easy.

( $\Rightarrow$ ) If  $G$  is 1-uniform, then  $\alpha(G) = 1$ .

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- *$G$  is 1-uniform if and only if  $G$  is a complete graph*
- *$G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs*
- *$G$  is 3-uniform if and only if  $G$  is the disjoint union of a 1-uniform and a 2-uniform graph.*

An *independent set* is a set of vertices in a graph, no two of which are adjacent.

$\alpha(G)$  denotes the maximum size of an independent set.

### Proof.

$(\Leftarrow)$  Easy.

$(\Rightarrow)$  If  $G$  is 1-uniform, then  $\alpha(G) = 1$ .

If  $G$  is 2-uniform, then  $|V(G)| \geq 2$  and  $G$  is  $P_4$ -free.

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

### Theorem

*If  $G$  is a graph, then*

- *$G$  is 1-uniform if and only if  $G$  is a complete graph*
- *$G$  is 2-uniform if and only if its complement  $\overline{G}$  is the disjoint union of one or more complete bipartite graphs*
- *$G$  is 3-uniform if and only if  $G$  is the disjoint union of a 1-uniform and a 2-uniform graph.*

An *independent set* is a set of vertices in a graph, no two of which are adjacent.

$\alpha(G)$  denotes the maximum size of an independent set.

### Proof.

$(\Leftarrow)$  Easy.

$(\Rightarrow)$  If  $G$  is 1-uniform, then  $\alpha(G) = 1$ .

If  $G$  is 2-uniform, then  $|V(G)| \geq 2$  and  $G$  is  $P_4$ -free.

Suppose that  $G$  is disconnected ...  $\overline{G}$  is  $K_{r,s}$





## Characterization of $k$ -Uniform Graphs for $k \leq 3$

Suppose that  $G$  is connected. ....

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

Suppose that  $G$  is connected. ....

Now suppose that  $G$  is 3-uniform. Recall that we may assume that  $G$  is twin-free.

*Claim:* Every two distinct vertices of  $G$  have exactly one common non-neighbor.

*Proof of Claim:* ...

## Characterization of $k$ -Uniform Graphs for $k \leq 3$

Suppose that  $G$  is connected. ....

Now suppose that  $G$  is 3-uniform. Recall that we may assume that  $G$  is twin-free.

*Claim:* Every two distinct vertices of  $G$  have exactly one common non-neighbor.

*Proof of Claim:* ...

Therefore, every two distinct vertices have a unique common neighbor in  $\overline{G}$ . By a theorem of Erdős,  $\overline{G}$  is isomorphic to a *friendship graph*, that is, a graph obtained from the disjoint union of  $n$  copies of  $K_2$  by adding to it a universal vertex.

By adding twins to the friendship graph, we see that the theorem is true.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

We will review the following work:

A.Erey, "Uniform length dominating sequence graphs", *Graphs and Combinatorics*, vol.36, pp.1819-1825, 2020.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

### Lemma

*Let  $G$  be a  $k$ -uniform graph and  $v$  be any vertex of  $G$ . Then,*

- *the subgraph  $G \setminus N[v]$  is  $(k - 1)$ -uniform,*
- *if  $G$  has no true twins, then  $G \setminus N[v]$  has no true twins.*

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

### Lemma

*Let  $G$  be a  $k$ -uniform graph and  $v$  be any vertex of  $G$ . Then,*

- *the subgraph  $G \setminus N[v]$  is  $(k - 1)$ -uniform,*
- *if  $G$  has no true twins, then  $G \setminus N[v]$  has no true twins.*

### Proof.

- ....

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

### Lemma

*Let  $G$  be a  $k$ -uniform graph and  $v$  be any vertex of  $G$ . Then,*

- *the subgraph  $G \setminus N[v]$  is  $(k - 1)$ -uniform,*
- *if  $G$  has no true twins, then  $G \setminus N[v]$  has no true twins.*

### Proof.

- ....
- Suppose that  $w_1, w_2$  are true twins of  $G \setminus N[v]$ ....



## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

### Lemma

*Let  $G$  be a  $k$ -uniform graph and  $v$  be any vertex of  $G$ . Then,*

- *the subgraph  $G \setminus N[v]$  is  $(k - 1)$ -uniform,*
- *if  $G$  has no true twins, then  $G \setminus N[v]$  has no true twins.*

### Proof.

- ....
- Suppose that  $w_1, w_2$  are true twins of  $G \setminus N[v]$ ....

$(w_2, w_3, \dots, w_k)$  is a legal dominating sequence of  $G \setminus N[v]$ ....

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

The author extended this result for  $k \geq 4$  as follows:

### Lemma

*Let  $G$  be a  $k$ -uniform graph and  $v$  be any vertex of  $G$ . Then,*

- *the subgraph  $G \setminus N[v]$  is  $(k - 1)$ -uniform,*
- *if  $G$  has no true twins, then  $G \setminus N[v]$  has no true twins.*

### Proof.

- ....
- Suppose that  $w_1, w_2$  are true twins of  $G \setminus N[v]$ ....

*$(w_2, w_3, \dots, w_k)$  is a legal dominating sequence of  $G \setminus N[v]$ ....*

*$(w_1, w_2, w_3, \dots, w_k, v)$  is also a legal dominating sequence of length  $(k + 1)$ , contradiction.*



## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Remark

*Let  $G$  be a graph with connected components  $G_1, \dots, G_c$ . Then  $G$  is  $k$ -uniform if and only if each  $G_i$  is  $k_i$ -uniform, where  $k = k_1 + \dots, k_c$  and  $k_i \geq 1$ .*

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Remark

*Let  $G$  be a graph with connected components  $G_1, \dots, G_c$ . Then  $G$  is  $k$ -uniform if and only if each  $G_i$  is  $k_i$ -uniform, where  $k = k_1 + \dots, k_c$  and  $k_i \geq 1$ .*

The *join* of two graphs  $G$  and  $H$ , denoted by  $G \vee H$  has vertex set  $V(G \vee H) = V(G) \cup V(H)$  and edge set  $E(G \vee H) = E(G) \cup E(H) \cup \{uv \mid u \in V(G), v \in V(H)\}$ .

# Characterization of $k$ -Uniform Graphs for $k \geq 4$

## Remark

*Let  $G$  be a graph with connected components  $G_1, \dots, G_c$ . Then  $G$  is  $k$ -uniform if and only if each  $G_i$  is  $k_i$ -uniform, where  $k = k_1 + \dots, k_c$  and  $k_i \geq 1$ .*

The *join* of two graphs  $G$  and  $H$ , denoted by  $G \vee H$  has vertex set  $V(G \vee H) = V(G) \cup V(H)$  and edge set  $E(G \vee H) = E(G) \cup E(H) \cup \{uv \mid u \in V(G), v \in V(H)\}$ .

Let  $\vee_t H$  denote the join of  $t$  copies of the graph  $H$  and  $tH$  denote disjoint union of  $t$  copies of  $H$

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Theorem

*If  $G$  is a  $k$ -uniform graph with  $k \geq 3$  and  $G$  has no true twins, then  $G$  is a disjoint union of 1-uniform and 2-uniform graphs.*

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Theorem

*If  $G$  is a  $k$ -uniform graph with  $k \geq 3$  and  $G$  has no true twins, then  $G$  is a disjoint union of 1-uniform and 2-uniform graphs.*

### Proof.

Strong induction on  $k$ . It has already been proved for  $k \leq 3$ . Let  $G$  be a graph with no true twins where  $k \geq 4$  and let  $v$  be a vertex of  $G$  ...

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Theorem

*If  $G$  is a  $k$ -uniform graph with  $k \geq 3$  and  $G$  has no true twins, then  $G$  is a disjoint union of 1-uniform and 2-uniform graphs.*

### Proof.

Strong induction on  $k$ . It has already been proved for  $k \leq 3$ . Let  $G$  be a graph with no true twins where  $k \geq 4$  and let  $v$  be a vertex of  $G$  ...

Let  $G_1, \dots, G_r$  be the 1-uniform connected components and  $H_1, \dots, H_t$  be the 2-uniform connected components of the subgraph  $G \setminus N[v]$ . ....



## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Theorem

*If  $G$  is a  $k$ -uniform graph with  $k \geq 3$  and  $G$  has no true twins, then  $G$  is a disjoint union of 1-uniform and 2-uniform graphs.*

### Proof.

Strong induction on  $k$ . It has already been proved for  $k \leq 3$ . Let  $G$  be a graph with no true twins where  $k \geq 4$  and let  $v$  be a vertex of  $G$  ...

Let  $G_1, \dots, G_r$  be the 1-uniform connected components and  $H_1, \dots, H_t$  be the 2-uniform connected components of the subgraph  $G \setminus N[v]$ . ....

Let us show that  $G$  is disconnected. Suppose to the contrary that  $G$  is connected. Let  $A_i = N(v_i) \cap N(v)$  and  $B_i = N(V(H_i)) \cap N(v)$  for each  $i$ .

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

### Theorem

*If  $G$  is a  $k$ -uniform graph with  $k \geq 3$  and  $G$  has no true twins, then  $G$  is a disjoint union of 1-uniform and 2-uniform graphs.*

### Proof.

Strong induction on  $k$ . It has already been proved for  $k \leq 3$ . Let  $G$  be a graph with no true twins where  $k \geq 4$  and let  $v$  be a vertex of  $G$  ...

Let  $G_1, \dots, G_r$  be the 1-uniform connected components and  $H_1, \dots, H_t$  be the 2-uniform connected components of the subgraph  $G \setminus N[v]$ . ....

Let us show that  $G$  is disconnected. Suppose to the contrary that  $G$  is connected. Let  $A_i = N(v_i) \cap N(v)$  and  $B_i = N(V(H_i)) \cap N(v)$  for each  $i$ .

Since  $G$  is connected,  $A_i$  and  $B_i$  are nonempty for each  $i$ . □

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ .

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ . We obtain a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ . We obtain a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $B_i \cap B_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $u_1$  and  $u_2$ .

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ . We obtain a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $B_i \cap B_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $u_1$  and  $u_2$ . Then  $(v, w, u'_1, u'_2, u_3, u'_3, \dots, v_1, \dots, v_r)$  is a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ . We obtain a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $B_i \cap B_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $u_1$  and  $u_2$ . Then  $(v, w, u'_1, u'_2, u_3, u'_3, \dots, v_1, \dots, v_r)$  is a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $A_i \cap B_j = \emptyset$  for every  $i$  and  $j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $u_1$ . Then again, a legal dominating sequence of length  $k - 1$  is found.



## Characterization of $k$ -Uniform Graphs for $k \geq 4$

Let us show that  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

- $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $v_2$ . We obtain a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $B_i \cap B_j = \emptyset$  whenever  $i \neq j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $u_1$  and  $u_2$ . Then  $(v, w, u'_1, u'_2, u_3, u'_3, \dots, v_1, \dots, v_r)$  is a legal dominating sequence of length  $r + 2t = k - 1$ , contradiction.
- $A_i \cap B_j = \emptyset$  for every  $i$  and  $j$ : Suppose that there exists a vertex  $w \in N(v)$  such that  $w$  is adjacent to both  $v_1$  and  $u_1$ . Then again, a legal dominating sequence of length  $k - 1$  is found.

Therefore,  $A_1, \dots, A_r, B_1, \dots, B_t$  are mutually disjoint.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

If  $r \geq 1$ , we consider removing the vertex  $v_1$  from  $G$ .

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

If  $r \geq 1$ , we consider removing the vertex  $v_1$  from  $G$ .

Notice that  $G \setminus N[v_1]$  is connected. However,  $G \setminus N[v_1]$  is  $(k - 1)$ -uniform and must be disconnected by the induction hypothesis.

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

If  $r \geq 1$ , we consider removing the vertex  $v_1$  from  $G$ .

Notice that  $G \setminus N[v_1]$  is connected. However,  $G \setminus N[v_1]$  is  $(k-1)$ -uniform and must be disconnected by the induction hypothesis.

If  $r = 0$ , then  $G \setminus N[v]$  has  $t$  connected 2-uniform components. So,  $k-1 = 2t$  and  $k \geq 5$ .

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

If  $r \geq 1$ , we consider removing the vertex  $v_1$  from  $G$ .

Notice that  $G \setminus N[v_1]$  is connected. However,  $G \setminus N[v_1]$  is  $(k-1)$ -uniform and must be disconnected by the induction hypothesis.

If  $r = 0$ , then  $G \setminus N[v]$  has  $t$  connected 2-uniform components. So,  $k-1 = 2t$  and  $k \geq 5$ .

First, let us show that  $u_1$  and  $u'_1$  have the same neighbors in  $N(v)$ ....

## Characterization of $k$ -Uniform Graphs for $k \geq 4$

If  $r \geq 1$ , we consider removing the vertex  $v_1$  from  $G$ .

Notice that  $G \setminus N[v_1]$  is connected. However,  $G \setminus N[v_1]$  is  $(k-1)$ -uniform and must be disconnected by the induction hypothesis.

If  $r = 0$ , then  $G \setminus N[v]$  has  $t$  connected 2-uniform components. So,  $k-1 = 2t$  and  $k \geq 5$ .

First, let us show that  $u_1$  and  $u'_1$  have the same neighbors in  $N(v)$ ....

We then show that  $G$  is disconnected and the result follows.