

# Domination Parameters in Graphs

Joanna Raczek

Faculty of Electronics, Telecommunications and Electronics  
Gdansk University of Technology

June, 2023

# Definitions

## Roman domination

### Roman dominating function


- Cockayne, Dreyer, Hedetniemi, and Hedetniemi in 2004 defined a *Roman dominating function* (RDF) on a graph  $G = (V, E)$  to be a function  $f: V \rightarrow \{0, 1, 2\}$  satisfying the condition that every vertex  $u$  for which  $f(u) = 0$  is adjacent to at least one vertex  $v$  for which  $f(v) = 2$ . [Roman domination in graphs](#), *Discrete Math.* **278** (2004).

# Definitions

## Roman domination

### Roman dominating function

- Cockayne, Dreyer, Hedetniemi, and Hedetniemi in 2004 defined a *Roman dominating function* (RDF) on a graph  $G = (V, E)$  to be a function  $f: V \rightarrow \{0, 1, 2\}$  satisfying the condition that every vertex  $u$  for which  $f(u) = 0$  is adjacent to at least one vertex  $v$  for which  $f(v) = 2$ . [Roman domination in graphs](#), *Discrete Math.* **278** (2004).




# Definitions

## Roman domination

### Roman dominating function

- Cockayne, Dreyer, Hedetniemi, and Hedetniemi in 2004 defined a *Roman dominating function* (RDF) on a graph  $G = (V, E)$  to be a function  $f: V \rightarrow \{0, 1, 2\}$  satisfying the condition that every vertex  $u$  for which  $f(u) = 0$  is adjacent to at least one vertex  $v$  for which  $f(v) = 2$ . [Roman domination in graphs, \*Discrete Math.\* \*\*278\*\* \(2004\).](#)



# Definitions

## Roman domination

### Roman dominating function

- ▶ For a real-valued function,  $f: V \rightarrow \mathbb{R}$ , the *weight* of  $f$  is  $w(f) = \sum_{v \in V} f(v)$ .
- ▶ The *Roman domination number*, denoted  $\gamma_R(G)$ , is the minimum weight of an RDF in  $G$ ; that is,  $\gamma_R(G) = \min\{w(f) : f \text{ is a RDF in } G\}$ .

# Definitions

## Roman domination

### Roman dominating function

- ▶ For a real-valued function,  $f: V \rightarrow \mathbb{R}$ , the *weight* of  $f$  is  $w(f) = \sum_{v \in V} f(v)$ .
- ▶ The *Roman domination number*, denoted  $\gamma_R(G)$ , is the minimum weight of an RDF in  $G$ ; that is,  $\gamma_R(G) = \min\{w(f) : f \text{ is a RDF in } G\}$ .




Figure:  $w(f) = 6$

# Definitions

## Roman domination

### Roman dominating function

- ▶ For a real-valued function,  $f: V \rightarrow \mathbb{R}$ , the *weight* of  $f$  is  $w(f) = \sum_{v \in V} f(v)$ .
- ▶ The *Roman domination number*, denoted  $\gamma_R(G)$ , is the minimum weight of an RDF in  $G$ ; that is,  $\gamma_R(G) = \min\{w(f) : f \text{ is a RDF in } G\}$ .




Figure:  $w(f) = 6$

# Definitions

## Roman domination

### Roman dominating function

- ▶ For a real-valued function,  $f: V \rightarrow \mathbb{R}$ , the *weight* of  $f$  is  $w(f) = \sum_{v \in V} f(v)$ .
- ▶ The *Roman domination number*, denoted  $\gamma_R(G)$ , is the minimum weight of an RDF in  $G$ ; that is,  $\gamma_R(G) = \min\{w(f) : f \text{ is a RDF in } G\}$ .




Figure:  $w(f) = 4$ ,  $\gamma_R(G) = 4$



# Historical background

## Roman domination

### Defending Roman Empire

- ▶ At the beginning of the 4th century AD, Roman Empire was consolidated and reformed by Emperor Constantine I (306–337).
- ▶ He did some remarkable things:
  - ▶ reorganized the economy by introducing golden coin solidus (or denarius),
  - ▶ allowing religious tolerance (Milan Edict in 312),
  - ▶ built new capital (Constantinople),
  - ▶ reorganized army.
- ▶ Constantine I placed armed forces not only on the borders but throughout all of the Empire.
- ▶ Major problem was payment of the army (which was up to 3/4 of the tax revenues), so optimal disposition was of utmost importance.

# Historical background

## Roman domination

### Defending Roman Empire

- ▶ Constantine I decided to differentiate communities of the Empire in three categories:
  - ▶ those with mobile and stationary troops,
  - ▶ those with only stationary troops,
  - ▶ those without troops at all.
- ▶ The condition was that communities without troops must be in neighborhood of communities with mobile troops, so that in case of attack they can be defended.

# Historical background

## Roman domination

### Defending Roman Empire

- ▶ Definition of a Roman dominating function was motivated by an article by I. Stewart, [Defend the Roman Empire!](#), *Scientific American*, 1999.
- ▶ Each vertex in graph represents a location in the Roman Empire.
- ▶ A location (vertex  $v$ ) is considered *unsecured* if no legions are stationed there (i.e.,  $f(v) = 0$ ) and *secured* otherwise (i.e., if  $f(v) \in \{1, 2\}$ ).
- ▶ An unsecured location (vertex  $v$ ) can be secured by sending a legion to  $v$  from an adjacent location (an adjacent vertex  $u$ ) with two legions.

# Why is it domination?

Roman domination

## Dominating set

- ▶ A set  $D \subseteq V$  is a *dominating set* of  $G$  if each vertex of  $V - D$  is adjacent to a vertex of  $D$ .
- ▶ The size of a smallest dominating set of  $G$  is the *domination number* of  $G$  and is denoted by  $\gamma(G)$ .

# Properties of Roman domination

Proposition (Cockayne et al., Roman domination in graphs (2004))

For any graph  $G$ ,

$$\gamma(G) \leq \gamma_R(G) \leq 2\gamma(G).$$

# Properties of Roman domination

Proposition (Cockayne et al., Roman domination in graphs (2004))

For any graph  $G$ ,

$$\gamma(G) \leq \gamma_R(G) \leq 2\gamma(G).$$

Proposition (Cockayne et al., Roman domination in graphs (2004))

For any graph  $G$ ,  $\gamma(G) = \gamma_R(G)$  if and only if  $G = \overline{K_n}$ .

# Properties of Roman domination

Proposition (Cockayne et al., Roman domination in graphs (2004))

For any graph  $G$ ,

$$\gamma(G) \leq \gamma_R(G) \leq 2\gamma(G).$$

Proposition (Cockayne et al., Roman domination in graphs (2004))

For any graph  $G$ ,  $\gamma(G) = \gamma_R(G)$  if and only if  $G = \overline{K_n}$ .

## Roman graphs

- ▶ Graphs which have Roman domination number equal to twice their domination number are called *Roman graphs*.

# Properties of Roman domination

Proposition (Cockayne et al., Roman domination in graphs (2004))

*Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then*



# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

*Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then*

- ▶  *$G[V_1]$  has maximum degree 1.*

# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

*Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then*

- ▶  *$G[V_1]$  has maximum degree 1.*
- ▶ *No edge of  $G$  joins  $V_2$  and  $V_1$ .*

# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then

- ▶  $G[V_1]$  has maximum degree 1.
- ▶ No edge of  $G$  joins  $V_2$  and  $V_1$ .
- ▶ Each vertex of  $V_0$  is adjacent to at most two vertices of  $V_1$ .

# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

*Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then*

- ▶  *$G[V_1]$  has maximum degree 1.*
- ▶ *No edge of  $G$  joins  $V_2$  and  $V_1$ .*
- ▶ *Each vertex of  $V_0$  is adjacent to at most two vertices of  $V_1$ .*
- ▶  *$V_2$  is a  $\gamma$ -set of  $G[V_0 \cup V_2]$*

# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then

- ▶  $G[V_1]$  has maximum degree 1.
- ▶ No edge of  $G$  joins  $V_2$  and  $V_1$ .
- ▶ Each vertex of  $V_0$  is adjacent to at most two vertices of  $V_1$ .
- ▶  $V_2$  is a  $\gamma$ -set of  $G[V_0 \cup V_2]$
- ▶ Let  $H = G[V_0 \cup V_2]$ . Then each vertex  $v \in V_2$  has at least two private neighbors relative to  $V_2$  in  $H$ .

# Properties of Roman domination

## Proposition (Cockayne et al., Roman domination in graphs (2004))

Let  $f = (V_0, V_1, V_2)$  be any  $\gamma_R$ -function. Then

- ▶  $G[V_1]$  has maximum degree 1.
- ▶ No edge of  $G$  joins  $V_2$  and  $V_1$ .
- ▶ Each vertex of  $V_0$  is adjacent to at most two vertices of  $V_1$ .
- ▶  $V_2$  is a  $\gamma$ -set of  $G[V_0 \cup V_2]$
- ▶ Let  $H = G[V_0 \cup V_2]$ . Then each vertex  $v \in V_2$  has at least two private neighbors relative to  $V_2$  in  $H$ .

Menti 1.

# Definitions

## Subgraph weakly induced

- ▶ Let  $G = (V, E)$  be a connected graph and let  $S \subseteq V$ .
- ▶ The *subgraph weakly induced* by  $S$  is the graph  $\langle S \rangle_w = (N[S], E_w)$ , where  $E_w$  consists of all edges in  $E$  having at least one vertex in  $S$ .

# Definitions

## Subgraph weakly induced

- ▶ Let  $G = (V, E)$  be a connected graph and let  $S \subseteq V$ .
- ▶ The *subgraph weakly induced* by  $S$  is the graph  $\langle S \rangle_w = (N[S], E_w)$ , where  $E_w$  consists of all edges in  $E$  having at least one vertex in  $S$ .




Figure: Graph  $G$



# Definitions

## Subgraph weakly induced

- ▶ Let  $G = (V, E)$  be a connected graph and let  $S \subseteq V$ .
- ▶ The *subgraph weakly induced* by  $S$  is the graph  $\langle S \rangle_w = (N[S], E_w)$ , where  $E_w$  consists of all edges in  $E$  having at least one vertex in  $S$ .




Figure: Graph  $\langle S \rangle_w$

# Definitions

## Weakly connected set

- ▶ A set  $S$  is a weakly connected set if  $\langle S \rangle_w$  is connected.

# Definitions

## Weakly connected set

- ▶ A set  $S$  is a weakly connected set if  $\langle S \rangle_w$  is connected.




Figure: Graph  $\langle S \rangle_w$  is not connected

# Definitions

## Weakly connected set

- ▶ A set  $S$  is a weakly connected set if  $\langle S \rangle_w$  is connected.




Figure: Graph  $\langle S \rangle_w$  is connected

# Definitions


## Weakly connected domination

- ▶ A set  $D \subseteq V(G)$  is a *dominating set* of  $G$  if every vertex  $v \in V(G) - D$  is adjacent to a vertex in  $D$ .
- ▶ A set  $D \subseteq V(G)$  is a *weakly connected dominating set* (WCDS) of  $G$  if  $D$  is dominating and  $\langle D \rangle_w$  is connected.
- ▶ The *weakly connected domination number* of  $G$ , denoted  $\gamma_{wc}(G)$ , is the minimum cardinality of a WCDS.

# Definitions

## Weakly connected domination

- ▶ A set  $D \subseteq V(G)$  is a *dominating set* of  $G$  if every vertex  $v \in V(G) - D$  is adjacent to a vertex in  $D$ .
- ▶ A set  $D \subseteq V(G)$  is a *weakly connected dominating set* (WCDS) of  $G$  if  $D$  is dominating and  $\langle D \rangle_w$  is connected.
- ▶ The *weakly connected domination number* of  $G$ , denoted  $\gamma_{wc}(G)$ , is the minimum cardinality of a WCDS.




# Definitions

## Weakly connected Roman domination

### Motivation

- We explore the idea of strengthening security of the Roman Empire by providing a better communication in emergency between the legions, while still having substantial costs of maintaining legions as low as possible.




# Definitions

## Weakly connected Roman domination

### Motivation

- We explore the idea of strengthening security of the Roman Empire by providing a better communication in emergency between the legions, while still having substantial costs of maintaining legions as low as possible.



*The single biggest problem in communication is the illusion that it has taken place.*

George Bernard Shaw



# Definitions

## Weakly connected Roman domination

### Motivation

- ▶ Two legions at different location (vertices  $u$  and  $v$ ) can *contact directly* if there is at most one unsecured location between them and the distance between  $u$  and  $v$  is at most 2 and they can *contact indirectly* if there is a sequence of vertices  $(u = u_1, u_2, \dots, u_k = v)$  such that  $u_i$  and  $u_{i+1}$  can contact directly for  $i = 1, 2, \dots, k - 1$ .

# Definitions

## Weakly connected Roman domination

### Motivation

- ▶ Two legions at different location (vertices  $u$  and  $v$ ) can *contact directly* if there is at most one unsecured location between them and the distance between  $u$  and  $v$  is at most 2 and they can *contact indirectly* if there is a sequence of vertices  $(u = u_1, u_2, \dots, u_k = v)$  such that  $u_i$  and  $u_{i+1}$  can contact directly for  $i = 1, 2, \dots, k - 1$ .
- ▶ The Roman Empire is *communicated* if any two legions at different locations can contact directly or indirectly.

# Definitions

## Weakly connected Roman domination

### Motivation

- ▶ A location of a legion is called *uncommunicated* if there exists another location of a legion such that the legions cannot contact directly nor indirectly.

# Definitions

## Weakly connected Roman domination

### Motivation

- ▶ A location of a legion is called *uncommunicated* if there exists another location of a legion such that the legions cannot contact directly nor indirectly.
- ▶ If the locations are uncommunicated, they cannot quickly inform the other locations nor ask them for help in case of urgent emergency.

# Definitions

## Weakly connected Roman domination

### Motivation

- ▶ A location of a legion is called *uncommunicated* if there exists another location of a legion such that the legions cannot contact directly nor indirectly.
- ▶ If the locations are uncommunicated, they cannot quickly inform the other locations nor ask them for help in case of urgent emergency.
- ▶ When all locations of legions are communicated, one can defend the Roman Empire more efficiently: supervise whole Empire and send orders to legions in reasonable time.

# Definitions

## Weakly connected Roman domination

### Weakly connected Roman dominating function


- We call the function  $f$  a *weakly connected Roman dominating function* in  $G$  (WCRDF) if each vertex  $u \in V_0$  is adjacent to a vertex  $v \in V_2$  and the subgraph  $\langle V_1 \cup V_2 \rangle_w$  weakly induced by  $V_1 \cup V_2$  is connected in  $G$ .

# Definitions

## Weakly connected Roman domination

### Weakly connected Roman dominating function

- We call the function  $f$  a *weakly connected Roman dominating function* in  $G$  (WCRDF) if each vertex  $u \in V_0$  is adjacent to a vertex  $v \in V_2$  and the subgraph  $\langle V_1 \cup V_2 \rangle_w$  weakly induced by  $V_1 \cup V_2$  is connected in  $G$ .



# Definitions

## Weakly connected Roman domination

### Weakly connected Roman dominating number

- ▶ We define the *weight*  $w(f)$  of  $f$  to be  $|V_1| + 2|V_2|$ .
- ▶ The *weakly connected Roman domination number*, denoted  $\gamma_R^{wc}(G)$ , is the minimum weight of a WCRDF in  $G$ .
- ▶ A WCRDF of weight  $\gamma_R^{wc}(G)$  we call a  $\gamma_R^{wc}(G)$ -function.



# Definitions

## Weakly connected Roman domination

### Weakly connected Roman dominating number

- ▶ We define the *weight*  $w(f)$  of  $f$  to be  $|V_1| + 2|V_2|$ .
- ▶ The *weakly connected Roman domination number*, denoted  $\gamma_R^{wc}(G)$ , is the minimum weight of a WCRDF in  $G$ .
- ▶ A WCRDF of weight  $\gamma_R^{wc}(G)$  we call a  $\gamma_R^{wc}(G)$ -function.




Figure:  $w(f) = 6$

# Definitions

## Weakly connected Roman domination

### Weakly connected Roman dominating number

- ▶ We define the *weight*  $w(f)$  of  $f$  to be  $|V_1| + 2|V_2|$ .
- ▶ The *weakly connected Roman domination number*, denoted  $\gamma_R^{wc}(G)$ , is the minimum weight of a WCRDF in  $G$ .
- ▶ A WCRDF of weight  $\gamma_R^{wc}(G)$  we call a  $\gamma_R^{wc}(G)$ -function.




Figure:  $w(f) = 5, \gamma_R^{wc}(G) = 5$

# Bounds

## Proposition

*If  $G$  is a connected graph, then*

$$\gamma_{wc}(G) \leq \gamma_R^{wc}(G) \leq 2\gamma_{wc}(G).$$

*The equality in the lower bound holds if and only if  $G = K_1$ .*

# Bounds

## Proposition

*For any connected graph  $G$  of order  $n$ ,*

$$\gamma_R^{\text{wc}}(G) \leq n.$$

*The equality  $\gamma_R^{\text{wc}}(G) = n$  holds if and only if  $G \in \{K_1, K_2\}$ .*

# Bounds

## Proposition

*If  $G = (V, E)$  is a connected graph of order at least 3 and  $f = (V_0, V_1, V_2)$  is a  $\gamma_R(G)$ -function, then*


$$\gamma_R(G) \leq \gamma_R^{wc}(G) \leq \gamma_R(G) + |V_2| - 1.$$

# Bounds

## Proposition

If  $G = (V, E)$  is a connected graph of order at least 3 and  $f = (V_0, V_1, V_2)$  is a  $\gamma_R(G)$ -function, then

$$\gamma_R(G) \leq \gamma_R^{wc}(G) \leq \gamma_R(G) + |V_2| - 1.$$




# Bounds

## Proposition

If  $G = (V, E)$  is a connected graph of order at least 3 and  $f = (V_0, V_1, V_2)$  is a  $\gamma_R(G)$ -function, then

$$\gamma_R(G) \leq \gamma_R^{wc}(G) \leq \gamma_R(G) + |V_2| - 1.$$




# Bounds

## Proposition

If  $G = (V, E)$  is a connected graph of order at least 3 and  $f = (V_0, V_1, V_2)$  is a  $\gamma_R(G)$ -function, then

$$\gamma_R(G) \leq \gamma_R^{wc}(G) \leq \gamma_R(G) + |V_2| - 1.$$



Menti 2.



# Certified dominating set


## Certified dominating set

- ▶ A subset  $S$  of  $V(G)$  is a *certified dominating set* of  $G$  if  $S$  is a dominating set and every vertex belonging to  $S$  has either zero or at least two neighbours in  $V(G) - S$ .
- ▶ M. Dettlaff, M. Lemańska, J. Topp, R. Ziemann, P. Żyliński, Certified domination, AKCE Int. J. of Graphs and Comb. (2018).

# Certified dominating set

## Certified dominating set


- ▶ A subset  $S$  of  $V(G)$  is a *certified dominating set* of  $G$  if  $S$  is a dominating set and every vertex belonging to  $S$  has either zero or at least two neighbours in  $V(G) - S$ .
- ▶ M. Dettlaff, M. Lemańska, J. Topp, R. Ziemann, P. Żyliński, Certified domination, AKCE Int. J. of Graphs and Comb. (2018).



# Certified domination number

## Certified domination number





- ▶ The cardinality of a minimum certified dominating set in  $G$  is called the *certified domination number* of  $G$  and is denoted  $\gamma_{\text{cer}}(G)$ .
- ▶ M. Dettlaff, M. Lemańska, J. Topp, R. Ziemann, P. Żyliński, Certified domination, AKCE Int. J. of Graphs and Comb. (2018).




# Certified dominating set

## Certified dominating set


- Every support vertex of a graph  $G$  belongs to every certified dominating set of  $G$ .




# Certified domination number



# Certified domination number



# Certified domination number



# Certified domination

## Theorem (Dettlaff et al. (2018))

*If  $G$  is a graph with no weak support, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*



# Certified domination

## Theorem (Dettlaff et al. (2018))

*If  $G$  is a graph with no weak support, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*

## Theorem (Dettlaff et al. (2018))

*Let  $G$  be a connected graph of order  $n$ . If  $G$  is the corona of some graph, then  $\gamma_{\text{cer}}(G) = n$ .*

# Certified domination

## Theorem (Dettlaff et al. (2018))

*If  $G$  is a graph with no weak support, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*

## Theorem (Dettlaff et al. (2018))

*Let  $G$  be a connected graph of order  $n$ . If  $G$  is the corona of some graph, then  $\gamma_{\text{cer}}(G) = n$ .*

## Theorem (Dettlaff et al. (2019))

*If a graph  $G$  has a unique  $\gamma(G)$ -set, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*

# Certified domination

## Theorem (Dettlaff et al. (2018))

*If  $G$  is a graph with no weak support, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*

## Theorem (Dettlaff et al. (2018))

*Let  $G$  be a connected graph of order  $n$ . If  $G$  is the corona of some graph, then  $\gamma_{\text{cer}}(G) = n$ .*

## Theorem (Dettlaff et al. (2019))

*If a graph  $G$  has a unique  $\gamma(G)$ -set, then  $\gamma(G) = \gamma_{\text{cer}}(G)$ .*

Menti 3

# Secondary Domination

## Motivation

- ▶ The secondary domination in graphs was defined by Hedetniemi, Rall et al., *Secondary domination in graphs*, AKCE Int. J. of Graphs and Comb. (2008).
- ▶ The authors of this paper mention a forthcoming paper on the algorithmic complexity of this domination problem, however this paper was never written.
- ▶ Since then some papers on this topic have appeared.

# Definitions

## $(1, 2)$ -domination

- ▶ A set  $D \subseteq V(G)$  is a  $(1, 2)$ -dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .

# Definitions

## $(1, 2)$ -domination

- ▶ A set  $D \subseteq V(G)$  is a  $(1, 2)$ -dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .
- ▶ The  $(1, 2)$ -domination number, denoted by  $\gamma_{1,2}(G)$ , is the cardinality of a smallest  $(1, 2)$ -dominating set of  $G$ .

# Definitions


## (1, 2)-domination

- ▶ A set  $D \subseteq V(G)$  is a (1, 2)-dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .
- ▶ The (1, 2)-domination number, denoted by  $\gamma_{1,2}(G)$ , is the cardinality of a smallest (1, 2)-dominating set of  $G$ .
- ▶ The (1, 2)-domination in graphs is a special case of (1,  $k$ )-domination. Here we only deal with the case when  $k = 2$ , that is, secondary domination.

# Definitions

## (1,2)-domination

- ▶ A set  $D \subseteq V(G)$  is a (1,2)-dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .
- ▶ The (1,2)-domination number, denoted by  $\gamma_{1,2}(G)$ , is the cardinality of a smallest (1,2)-dominating set of  $G$ .






# Definitions

## (1,2)-domination


- ▶ A set  $D \subseteq V(G)$  is a (1,2)-dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .
- ▶ The (1,2)-domination number, denoted by  $\gamma_{1,2}(G)$ , is the cardinality of a smallest (1,2)-dominating set of  $G$ .



# Definitions

## (1,2)-domination

- ▶ A set  $D \subseteq V(G)$  is a (1,2)-dominating set if each vertex  $v$  of  $V - D$  has a neighbour in  $D$  as well as another vertex of  $D$  at a distance not greater than 2 from  $v$ .
- ▶ The (1,2)-domination number, denoted by  $\gamma_{1,2}(G)$ , is the cardinality of a smallest (1,2)-dominating set of  $G$ .



# (1,2)-domination

## Theorem (Hedetniemi et al. (2008))

*If  $G$  is a graph, then  $\gamma(G) \leq \gamma_{1,2}(G) \leq \gamma_t(G)$ .*

# (1,2)-domination

## Theorem (Hedetniemi et al. (2008))

*If  $G$  is a graph, then  $\gamma(G) \leq \gamma_{1,2}(G) \leq \gamma_t(G)$ .*

## Theorem (JR.)

*If  $G$  is a graph with  $\delta(G) > 1$  and without a triangle, then*

$$\gamma(G) = \gamma_{1,2}(G).$$

# (1,2)-domination

## Theorem (Hedetniemi et al. (2008))

*If  $G$  is a graph, then  $\gamma(G) \leq \gamma_{1,2}(G) \leq \gamma_t(G)$ .*

## Theorem (JR.)

*If  $G$  is a graph with  $\delta(G) > 1$  and without a triangle, then*

$$\gamma(G) = \gamma_{1,2}(G).$$

Menti 4

# Thank you!